

AD-A141 512

SUBSTITUTION AND ADDITION REACTIONS OF A
(METHYLENE)PHOSPHINE WITH ALKYL... (U) TEXAS CHRISTIAN
UNIV FORT WORTH DEPT OF CHEMISTRY B L LI ET AL.


1/1

UNCLASSIFIED

18 MAY 84 TCU/DC/TR-84-04 N00014-79-C-0632 F/G 7/3

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

AD-A141 512

OFFICE OF NAVAL RESEARCH
Contract No. N00014-79-C-0632

Task No. NR-053-720

TECHNICAL REPORT NO. TCU/DC/TR-84-04

Substitution and Addition Reactions of a
(Methylene)phosphine with Alkylolithium Reagents

by

Bi-L. Li and R.H. Neilson

Prepared for Publication

in

Inorganic Chemistry

Texas Christian University
Chemistry Department
Fort Worth, TX 76129

May 18, 1984

DTIC
ELECTE
MAY 30 1984
S B

DTIC FILE COPY

Reproduction in whole or in part is
Permitted for any purpose of the United States Government

This document has been approved for public
release and sale; its distribution is unlimited

84 05 29 003

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TCU/DC/TR-84-04	2. GOVT ACCESSION NO. AD - A141512	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Substitution and Addition Reactions of a (methylene)phosphine with Alkylolithium Reagents		5. TYPE OF REPORT & PERIOD COVERED Technical Report, 1983
		6. PERFORMING ORG. REPORT NUMBER TR-84-04
7. AUTHOR(s) Bi-L. Li and R.H. Neilson		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0632
9. PERFORMING ORGANIZATION NAME AND ADDRESS Texas Christian University Fort Worth, Texas 76129		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-720
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research		12. REPORT DATE May 18, 1984
		13. NUMBER OF PAGES 10
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Prepared for Publication in <u>Inorganic Chemistry</u>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) (Methylene)phosphine [bis(trimethylsilyl)methyl]phosphine diphosphinomethane trialkylphosphine		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Treatment of the (methylene)phosphine $(\text{Me}_3\text{Si})_2\text{NP}=\text{CHSiMe}_3$ (1) with alkylolithium reagents and Me_3SiCl affords either the diphosphinomethane derivative $\text{R}_2\text{P}-\text{CH}(\text{SiMe}_3)-\text{P}(\text{R})\text{CH}(\text{SiMe}_3)_2$ (2, R = Me) or the trialkylphosphine $\text{R}_2\text{PCH}(\text{SiMe}_3)_2$ (3, R = <u>t</u> -Bu). A reaction pathway involving both nucleophilic substitution and addition reactions at the P=C moiety is suggested. Compound <u>2</u> exhibits hindered rotation about the P-CH(SiMe ₃) ₂ bond with a measured $\Delta G^\ddagger_{\text{C}}$ of 17.7 kcal/mole.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Substitution and Addition Reactions of
a (Methylene)phosphine with Alkylolithium Reagents

Bei-Li Li and Robert H. Neilson*

Department of Chemistry
Texas Christian University
Fort Worth, Texas 76129

Abstract

✓ Treatment of the (methylene)phosphine $(\text{Me}_3\text{Si})_2\text{NP}=\text{CHSiMe}_3$ (1) with alkylolithium reagents and Me_3SiCl affords either the diphosphinomethane derivative $\text{R}_2\text{P}-\text{CH}(\text{SiMe}_3)-\text{P}(\text{R})\text{CH}(\text{SiMe}_3)_2$ (2, $\text{R} = \text{Me}$) or the trialkylphosphine $\text{R}_2\text{PCH}(\text{SiMe}_3)_2$ (3, $\text{R} = t\text{-Bu}$). A reaction pathway involving both nucleophilic substitution and addition reactions at the $\text{P}=\text{C}$ moiety is suggested. Compound 2 exhibits hindered rotation about the $\text{P}-\text{CH}(\text{SiMe}_3)_2$ bond with a measured $\Delta G^\ddagger_{\text{C}}$ of 17.7 kcal/mole.

delta G[‡] sub C


Accession For	
NTIS GR&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	




Many recent reports have dealt with the preparative chemistry of (methylene)phosphines $RP=CR'_2$ as well as their structural and bonding characteristics.¹ Moreover, some relatively clear and consistent patterns of reactivity of such compounds are now emerging. The types of reactions of (methylene)phosphines reported thus far include: (1) addition of polar electrophilic reagents to the $P=C$ bond², (2) oxidation to 3-coordinate $P(V)$ derivatives³, (3) complexation of transition metals to the phosphorus lone pair or to the $P=C$ π bond⁴, and (4) various cycloaddition processes such as Diels-Alder reactions.⁵ In another general mode of reactivity, we find that these coordinatively unsaturated phosphorus compounds will also react smoothly with nucleophiles.⁶ We report here some novel examples involving both substitution and addition reactions of a bis(trimethylsilyl)amino substituted (methylene)phosphine with alkyllithium reagents.

Treatment of [bis(trimethylsilyl)amino](trimethylsilyl-methylene)phosphine^{2c} 1 in Et_2O at $-78^\circ C$ with $MeLi$, followed by quenching with Me_3SiCl , does not yield the expected⁷ phosphine $(Me_3Si)_2NP(Me)CH(SiMe_3)_2$. Instead, the reaction takes a much more complicated course, forming the novel diphosphinomethane derivative 2 (eq 1). The product 2 is isolated as a colorless liquid in 88% yield by fractional distillation. The characterization of 2 is based mainly on NMR data (Table I) with the AB pattern in the ^{31}P NMR spectrum being especially diagnostic. A satisfactory elemental analysis and the

appropriate mass spectral fragmentation pattern were also obtained for 2.



Under similar conditions, the reaction of 1 with t-BuLi occurs in a 1:2 stoichiometry (eq 2) to afford the di(t-butyl)-phosphine 3 in 69% yield rather than a P-C-P product analogous to 2.



These remarkably clean reactions of the (methylene)phosphine 1 with alkylolithium reagents must involve both nucleophilic displacement of the bis(trimethylsilyl)amide group as well as addition across the P=C bond. The reaction pathway proposed in Scheme I is strongly suggested by the following experimental observations: (1) The byproduct $(\text{Me}_3\text{Si})_3\text{N}$ can be isolated and identified by NMR, indicating that $(\text{Me}_3\text{Si})_2\text{NLi}$ is indeed formed.⁸

Moreover, the ^1H NMR integration of the crude reaction product is consistent with the stoichiometry given by equation 1. (2) The use of 1:1 mole ratios of 1 to RLi yields only unreacted 1 in addition to products 2 or 3. (3) The formation of 3 in the $t\text{-BuLi}$ reaction rather than a product similar to 2 is consistent with increased steric hindrance at phosphorus in intermediates A and B when $\text{R} = t\text{-Bu}$. (4) When the reaction with MeLi is monitored by low-temperature ^{31}P NMR, the (methylene)phosphine intermediate A is not seen but an AB pattern ($\delta_{\text{A}} -1.8$, $\delta_{\text{B}} -6.8$, $J_{\text{AB}} = 69.3$ Hz) assignable to the diphosphorus anion C is clearly observed prior to the addition of Me_3SiCl .

A final point of interest concerns the ^1H and ^{13}C NMR spectra of the di(t -butyl)phosphine 3 which reveal non-equivalence of the $t\text{-Bu}$ signals at room temperature. A high temperature ^1H NMR study shows that the two t -butyl doublets coalesce to a single doublet at 65°C (in C_6H_6 solution) indicating a barrier to rotation about the $\text{P-CH}(\text{SiMe}_3)_2$ bond of approximately 17.7 kcal/mole. The extreme steric hindrance caused by the bis(trimethylsilyl)methyl group in phosphorus (III) compounds has been previously noted.⁹

Experimental Section

Materials and General Procedures. The (methylene)phosphine 1 was prepared according to the published procedure.^{2c} Alkyl lithium reagents were used as received from commercial sources. Ether and hexane were distilled from CaH_2 prior to use.

Proton NMR spectra were recorded on a Varian EM-390 spectrometer; ^{13}C and ^{31}P , both with ^1H decoupling, were obtained on a JEOL FX-60 instrument. Mass spectral data were obtained on Finnigan OWA 1020 GC-MS system. Elemental analyses were performed by Schwarzkopf Microanalytical Laboratory, Woodside, N.Y.

All reactions and other manipulations were carried out under an atmosphere of dry nitrogen or under vacuum. The procedures described below are typical of those which gave the best yields of compounds 2 and 3. Preliminary experiments using equimolar quantities of 1 and the alkyllithium reagents afforded the same products together with unreacted (methylene)phosphine.

Reaction of 1 with MeLi. Methyllithium (30 mmol, 21.5 mL, 1.4 M in Et_2O) was added at -78°C to a stirred solution of 1 (20 mmol, 5.6 g) in Et_2O (50 mL) to yield a light yellow precipitate, presumably anion C (Scheme I). After stirring the mixture for 2.5 h at -78°C , Me_3SiCl (30 mmol, 3.8 mL) was added and the mixture was allowed to warm to room temperature. After stirring overnight, Et_2O was removed under vacuum and hexane (20 mL) was added. The mixture was then filtered and the solvent and $(\text{Me}_3\text{Si})_3\text{N}$ (identified by ^1H NMR) were removed under vacuum. Distillation through a short path column afforded compound 2 as a colorless liquid (3.1 g, 87% yield, bp $99-105^\circ\text{C}/0.05$ mm). Anal. Calcd: C, 47.73; H, 10.80. Found: C, 47.51; H, 10.77. Mass spectrum, m/e (relative intensity): 352 (2), 337 (9), 291 (22), 279 (6), 205 (4), 193 (10), 147 (14), 73 (100).

Reaction of 1 with t -BuLi. By means of a similar procedure, 1 (20 mmol) was treated with t -BuLi (40 mmol, 22.2 mL, 1.8 M in pentane) at -78°C in Et_2O (50 mL). After quenching with Me_3SiCl (40 mmol) and work-up as described above, distillation gave 3 as a colorless liquid (4.2 g, 69% yield, bp $82-83^{\circ}\text{C}/0.15$ mm). Anal. Calcd: C, 59.21; H, 12.17. Found: C, 59.08; H, 12.05. Mass spectrum, m/e (relative intensity): 304 (22), 289 (25), 248 (81), 159 (28), 145 (14), 73 (100).

Acknowledgment. We thank the U.S. Office of Naval Research and the Robert A. Welch Foundation for generous financial support.

References and Notes

- (1) For a general review see: Appel, R.; Knoll, F.; Ruppert, I. Angew. Chem. Int. Ed. Engl. 1981, 20, 731.
- (2) (a) Kleback, T.C.; Lourens, R.; Bickelhaupt, F. J. Am. Chem. Soc. 1978, 100, 4887.
- (b) Issleib, K.; Schmidt, H.; Wirkner, Ch. Z. Anorg. Allg. Chem. 1981, 473, 85.
- (c) Neilson, R.H. Inorg. Chem. 1981, 20, 1679.
- (3) (a) Niecke, E.; Wildbrecht, D.-A. J. Chem. Soc., Chem. Commun. 1981, 72.
- (b) Caira, M.; Neilson, R.H.; Watson, W.H.; Wisian-Neilson, P.; Xie, Z.-M. J. Chem. Soc., Chem. Commun., in press.
- (4) (a) Kroto, H.W.; Nixon, J.F.; Taylor, M.J.; Frew, A.A.; Muir, K. Polyhedron 1982, 1, 89.
- (b) van der Knaap, Th. A.; Bickelhaupt, F.; van der Poel, H.; van Koten, G.; Stam, C.H. J. Am. Chem. Soc. 1982, 104, 1756.
- (c) Cowley, A.H.; Jones, R.A.; Stewart, C.A.; Stuart, A.L. J. Am. Chem. Soc. 1983, 105, 3737.
- (d) Al-Resayes, S.I.; Klein, S.I.; Kroto, H.W.; Meidine, M.F.; Nixon, J.F. J. Chem. Soc., Chem. Commun. 1983, 930.
- (e) van der Knaap, Th.A.; Jenneskens, L.W.; Meenwissen, H.J.; Bickelhaupt, F.; Walther, D.; Dinjus, E.; Uhlig, E.; Spek, A.L. J. Organomet. Chem. 1983, 254, C33.

- (f) Neilson, R.H.; Thoma, R.J.; Vickovic, I.; Watson, W.H.
Organometallics, submitted for publication.
- (5) (a) Appel, R.; Korte, S.; Halstenberg, M.; Knach, F. Chem. Ber. 1982, 115, 3610.
(b) Meriem, A.; Majoral, J.-P.; Revel, M.; Navech, J.
Tetrahedron Lett. 1983, 24, 1975.
(c) van der Knaap, Th. A.; Bickelhaupt, F. Tetrahedron 1983, 38, 3189.
(d) Appel, R.; Zimmermann, R. Tetrahedron Lett. 1983, 24, 3591.
- (6) Nucleophilic displacement of chloride from P-Chloro substituted methylenephosphines is known:
(a) Appel, R.; Kundgen, U. Angew. Chem. Int. Ed. Engl. 1982, 21, 219.
(b) Cowley, A.H.; Ebsworth, E.A.V.; Kemp, R.A.; Rankin, D.W.H.; Stewart, C.A. Organometallics 1982, 1, 1720.
(c) Cowley, A.H.; Kilduff, J.E.; Pakulski, M.; Stewart, C.A. J. Am. Chem. Soc. 1983, 105, 1655.
- (7) Addition of MeLi to the analogous iminophosphine $(\text{Me}_3\text{Si})_2\text{NP}=\text{NSiMe}_3$ does yield the simple P-Me derivative:
Cowley, A.H.; Kemp, R.A.; J. Chem. Soc., Chem. Commun. 1982, 319.
- (8) Displacement of the $(\text{Me}_3\text{Si})_2\text{N}$ group from the analogous iminophosphine by very bulky organolithium compounds has recently been reported: Romanenko, V.D.; Ruben, A.V.; Markovski, L.N. J. Chem. Soc., Chem. Commun. 1983, 187.
- (9) Goldwhite, H.; Power, P.P. Org. Magn. Res. 1978, 11, 499.

Table I. NMR Spectroscopic Data^a

compd	Signal	¹ H		¹³ C		³¹ P
		δ	J _{PH}	δ	J _{PC}	δ
$\text{Me}_2\text{P}-\text{CHSiMe}_3$ $\quad $ $\text{Me}-\text{P}-\text{CH}(\text{SiMe}_3)_2$ $\quad \sim$ $\quad 2$	<u>Me₃SiC</u>	0.14	1.0	2.56	5.4,	-23.7 ^b ,
					1.8	-37.9
	(<u>Me₃Si</u>) ₂ C ^c	0.10	0.4	1.89	6.7	
				3.49	2.4	
	<u>Me₂PC^c</u>	1.15	4.8	<u>d</u>		
		1.16	4.2			
	<u>MeP</u>	1.30	6.0	<u>d</u>		
	<u>Si₂CH</u>	1.04	2.4	<u>d</u>		
	<u>P₂CH</u>	<u>e</u>		22.80	48.0,	
					33.3	
$(\text{t-Bu})_2\text{P}-\text{CH}(\text{SiMe}_3)_2$ $\quad \sim$ $\quad 3$	<u>Me₃SiC</u>	0.12	0.2	2.48	11.0	47.6
		0.18		5.12		
	CH	1.15	13.5	28.67	55.6	
	<u>Me₃C^c</u>	1.17	12.3	30.20	14.0	
		1.09	10.8	32.02	16.5	
	<u>Me₃C^c</u>			30.95	30.5	
				32.42	17.1	

^a Chemical shifts are downfield from Me₄Si for ¹H and ¹³C spectra and from H₃PO₄ for ³¹P spectra; coupling constants in Hz. Solvents: ¹H, CH₂Cl₂; ¹³C and ³¹P, CDCl₃.







^b AB pattern with J_{pp} = 95.2 Hz.

^c Diastereotopic groups observed in ¹H and/or ¹³C NMR spectra.

^d Multiplet of ca. 16 overlapping peaks in range δ 9.8 - δ 15.1.

^e Signals obscured by MeP resonances.

Scheme I



END

FILMED

1/11/71

DATE